
# CBSE PMT - 2007 SCREENING

## **PHYSICS**

- 1. The primary and secondary coil of a transformer have 50 and 1500 turns respectively. If the magnetic flux  $\phi$  linked with the primary coil is given by  $\phi = \phi_0 + 4t$ , where  $\phi$  is in webers, t is time in seconds and  $\phi_0$  is a constant, the output voltage across the secondary coil is
  - (a) 120 volts
- (b) 220 volts
- (c) 30 volts
- (d) 90 volts
- A beam of electron passes undeflected through mutually perpendicular electric and magnetic fields. If the electric field is switched off, and the same magnetic field is maintained, the electrons move
  - (a) in a circular orbit
  - (b) along a parabolic path
  - (c) along a straight line
  - (d) in an elliptical orbit.
- The position x of a particle with respect to time t along x-axis is given by  $x = 9t^2 - t^3$  where x is in metres and t in second. What will be the position of this particle when it achieves maximum speed along the +ve x direction?
  - (a) 54m (b) 81m (c) 24m (d) 32 m.
- A particle starting from the origin (0,0) moves in a straight line in the (x, y) plane. Its coordinates at a later time are  $(\sqrt{3},3)$ . The path of the particle makes with the x-axis an angle of
  - (d) 30° (a) 45° (b)  $60^{\circ}$  (c)  $0^{\circ}$
- A car moves from X to Y with a uniform speed v<sub>11</sub> and returns to Y with a uniform speed v<sub>d</sub>. The average speed for this round trip is
- (b)  $\frac{v_d v_u}{v_d + v_u}$
- (c)  $\frac{v_u + v_d}{2}$  (d)  $\frac{2v_d v_u}{v_d + v_u}$
- Two radioactive substances A and B have decay constants  $5\lambda$  and  $\lambda$  respectively. At t = 0 they have the same number of nuclei. The ratio of number of nuclei of A to those of B will be  $(1/e)^2$ after a time interval
  - (a)  $4\lambda$
- (b)  $2\lambda$  (c)  $1/2\lambda$  (d)  $1/4\lambda$

7. In the following circuit, the output Y for all possible inputs A and B is expressed by the truth table




- (a) A B Y (b) A B Y 0 1 1 001 0 1 1 0 1 0 101 100 1 1 0 1 1 0
- (c) A B Y (d) ABY  $0 \ 0 \ 0$ 000 0 1 1 0 1 0 101 100 1 1 1 1 1 1
- In a radioactive decay process, the negatively charged emitted  $\beta$ -particles are
- (a) the electrons produced as a result of the decay of neutrons inside the nucleus
- (b) the electrons produced as a result of collisions between atoms
- (c) the electrons orbiting around the nucleus
- (d) the electrons present inside the nucleus.
- The phase difference between the instantaneous velocity and acceleration of a particle executing simple harmonic motion is
  - (a)  $\pi$
- (b)  $0.707 \,\pi$
- (c) zero
- (d)  $0.5 \,\pi$
- In a mass spectrometer used for measuring the masses of ions, the ions are initially accelerated by an electric potential V and then made to describe semicircular path of radius R using a magnetic field B. If V and B are kept constant,

the ratio 
$$\left(\frac{\text{charge on the ion}}{\text{mass of the ion}}\right)$$
 will be


proportional to

- (a)  $1/R^2$ 
  - (b) R<sup>2</sup>
- (c) R
- (d) 1/R
- A wheel has angular acceleration of 3.0 rad/sec<sup>2</sup> and an initial angular speed of 2.00 rad/sec. In a time of 2 sec it has rotated through an angle (in radian) of
  - (a) 10
- (b) 12
- (c) 4
- (d) 6

12. A hollow cylinder has a charge q coulomb within i. If  $\phi$  is the electric flux in units of voltmeter associated with the curved surface B, the flux linked with the plane surface A in units of voltmeter will be



- (d)  $\frac{1}{2} \left( \frac{q}{\epsilon_0} \phi \right)$
- The frequency of a light wave in a material is  $2 \times 10^{14}$  Hz and wavelength is 5000 Å. The refractive index of material will be
  - (a) 1.50
- (b) 3.00 (c) 1.33
- (d) 1.40
- A nucleus  ${}_{Z}^{A}X$  has mass represented by M(A, Z). If  $M_p$  and  $M_n$  denote the mass of proton and neutron respectively and B.E. the binding energy in MeV, then
  - (a) B.E. =  $[ZM_p + (A-Z)M_p M(A, Z)]c^2$ (b) B.E. =  $[ZM_p + ZM_n M(A, Z)]c^2$ (c) B.E. =  $M(A, Z) ZM_p (A-Z)M_n$ (d) B.E. =  $M(A, Z) ZM_p (A-Z)M_n$
- The total power dissipated in watts in the circuit shown here is




- (a) 40
- (b) 54
- (c) 4

(d) 16

- 16. If the nucleus  $^{27}_{13}$ Al has nuclear radius of about
  - 3.6 fm, then  ${}^{125}_{32}$ Te would have its radius approximately as
  - (a) 9.6 fm
- (b) 12.0 fm
- (c) 4.8 fm
- (d) 6.0 fm.

A block B is pushed momentarily along a 17. horizontal surface with an initial velocity V. If  $\mu$ is the coefficient of sliding friction between B and the surface, block B will come to rest after a time



- (a)  $g\mu/V$
- (b) g/V
- (c) V/g
- (d)  $V/(g\mu)$ .
- A particle moving along x-axis has acceleration

f, at time t, given by 
$$f = f_0 \left( 1 - \frac{t}{T} \right)$$
, where  $f_0$ 

and T are constants. The particle at t=0 has zero velocity. In the time interval between t = 0and the instant when f = 0, the particle's velocity  $(v_{v})$  is

- (a)  $\frac{1}{2} f_0 T^2$  (b)  $f_0 T^2$
- (c)  $\frac{1}{2}f_0T$
- (d)  $f_0T$ .
- A common emitter amplifier has a voltage gain of 50, an input impedance of  $100\Omega$  and an output impedance of  $200\Omega$ . The power gain of the amplifier is
  - (a) 1000
- (b) 1250 (c) 100
- Three resistances P, Q, R each of  $2\Omega$  and an unknown resistance S form the four arms of a Wheatstone bridge circuit. When a resistance of  $6\Omega$  is connected in parallel to S the bridge


(d) 500.

- (a)  $3\Omega$
- (b)  $6\Omega$
- (c) 1Ω
- (d)  $2\Omega$
- A steady current of 1.5 amp flows through a copper voltameter for 10 minutes. If the electrochemical equivalent of copper is  $30 \times 10^{-5}$  g coulomb<sup>-1</sup>, the mass of copper deposited on the electrode will be

gets balanced. What is the value of S?

(a)  $0.50\,\mathrm{g}$  (b)  $0.67\,\mathrm{g}$  (c)  $0.27\,\mathrm{g}$  (d)  $0.40\,\mathrm{g}$ .


A small coin is resting on the bottom of a beaker filled with liquid. A ray of light from the coin travels upto the surface of the liquid and moves along its surface. How fast is the light travelling in the liquid?




- (a)  $2.4 \times 10^8 \,\text{m/s}$
- (b)  $3.0 \times 10^8 \,\text{m/s}$
- (c)  $1.2 \times 10^8 \,\text{m/s}$
- (d)  $1.8 \times 10^8 \,\text{m/s}$
- 23. A transformer is used to light a 100 W and 110 V lamp from a 220 V mains. If the main current is 0.5 amp, the efficiency of the transformer is approximately
  - (a) 50% (b) 90% (c) 10%
- Dimensions of resistance in an electrical circuit, in terms of dimension of mass M, of length L, of time T and of current I, would be
  - (a)  $ML^2T^{-2}$
- (b)  $ML^2T^{-1}I^{-1}$
- (c)  $ML^2T^{-3}I^{-2}$
- (d)  $ML^2T^{-3}I-1$
- A uniform rod AB of length  $\ell$ , and mass m is free to rotate about point A. The rod is released from rest in the horizontal position. Given that the

moment of inertia of the rod about A is  $\frac{m\ell^2}{3}$ ,

the initial angular acceleration of the rod will be



A particle of mass m moves in the XY plane with a velocity v along the straight line AB. If the angular momentum of the particle with respect tociginO isL<sub>A</sub> when it is at A and L<sub>B</sub> when it is




- (b) the relationship between  $L_A$  and  $L_B$  depends upon the slope of the line AB

- (d)  $L_A > L_B$ . The particle executing simple harmonic motion 27.

has a kinetic energy  $K_0 \cos^2 \omega t$ . The maximum values of the potential energy and the total energy are respectively

- (a)  $K_0/2$  and  $K_0$
- (c)  $K_0$  and  $K_0$
- (b) K<sub>0</sub> and 2K<sub>0</sub>(d) 0 and 2K<sub>0</sub>.
- Charges +q and -q are placed at points A and B respectively which are a distance 2L apart, C is the midpoint between A and B. The work done in moving a charge +Q along the semicircle CRD




- Monochromatic light of frequency  $6.0 \times 10^{14} \, \text{Hz}$ is produced by a laser. The power emitted is  $2 \times 10^{-3}$  w. The number of photons emitted, on the average, by the sources per second is the average (a)  $5 \times 10^{16}$  (b)  $5 \times 10^{15}$  (c)  $5 \times 10^{14}$  (d)  $5 \times 10^{15}$

- 30. Three point charges +q, -q and +q are placed at points (x = 0, y = a, z = 0), (x = 0, y = 0, z = 0) and (x = a, y = 0, z = 0) respectively. The magnitude and direction of the electric dipole moment vector of this charge assembly are
  - (a)  $\sqrt{2}$ qa along the line joining points (x = 0, y = 0, z = 0) and (x = a, y = a, z = 0)
  - (b) qa along the line joining points (x = 0, y = 0,z = 0) and (x = a, y = a, z = 0)
  - (c)  $\sqrt{2}$ qa along +ve x direction
  - (d)  $\sqrt{2}$ qa along +ve y direction
- Two satellites of earth,  $S_1$  and  $S_2$  are moving in the same orbit. The mass of  $S_1$  is four times the mass of S<sub>2</sub>. Which one of the following statements is true?
  - (a) The potential energies of earth satellites in the two cases are equal.
  - (b)  $S_1$  and  $S_2$  are moving with the same speed.
  - (c) The kinetic energies of the two satellites are
  - (d) The time period of  $S_1$  is four times that of  $S_2$ .
- $\overrightarrow{A}$  and  $\overrightarrow{B}$  are two vectors and  $\theta$  is the angle between them, if  $|\overrightarrow{A} \times \overrightarrow{B}| = \sqrt{3}(\overrightarrow{A}.\overrightarrow{B})$ , the value

of  $\theta$  is

- (a)  $45^{\circ}$ (b)  $30^{\circ}$  (c)  $90^{\circ}$ (d) 60°
- 33. A mass of 2.0 kg is put on a flat pan attached to a vertical spring fixed on the ground as shown in the figure. The mass of the spring and the pan is negligible. When pressed slightly and released the mass executes a simple harmonic motion. The spring constant is 200 N/m. What should be the minimum amplitude of the motion so that the mass gets detached from the pan (take g = 10 $m/s^2$ )?
  - (a) 10.0 cm
  - (b) any value less than 12.0 cm
  - (c) 4.0 cm
  - (d) 8.0 cm
- A charged paritcle (charge q) is moving in a circle of radius R with uniform speed v. The associated magnetic moment µ is given by
  - (a)  $qvR^2$
- (b)  $qvR^{2}/2$
- (c) qvR
- (d) qvR/2

- 35. The total energy of electron in the ground state of hydrogen atom is – 13.6 eV. The kinetic energy of an electron in the first excited state is
  - (a) 6.8 eV
- (b) 13.6eV
- (c) 1.7 eV
- (d) 3.4 eV.
- A 5 watt source emits monochromatic light of wavelength 5000 Å. When placed 0.5 m away, it liberates photoelectrons from a photosensitive metallic surface. When the source is moved to a distance of 1.0 m, the number of photoelectrons liberated will be reduced by a factor of
  - (a) 8
- (b) 16
- (c) 2
- (d) 4
- A particle executes simple harmonic oscillation with an amplitude a. The period of oscillation is T. The minimum time taken by the particle to travel half of the amplitude from the equilibrium position is
  - (a) T/8
- (b) T/12 (c) T/2
- (d) T/4
- For a cubic crystal structure which one of the following relations indicating the cell characteristics is correct?
  - (a)  $a \neq b \neq c$  and  $\alpha = \beta = \gamma = 90^{\circ}$
  - (b) a = b = c and  $\alpha \neq \beta \neq \gamma = 90^{\circ}$
  - (c) a = b = c and  $\alpha = \beta = \gamma = 90^{\circ}$
  - (c)  $a \neq b \neq c$  and  $\alpha \neq \beta$  and  $\gamma \neq 90^{\circ}$
- Two condensers, one of capacity C and other of capacity C/2 are connected to a V-volt battery, as shown.



The work done in charging fully both the condensers is

- (a)  $\frac{1}{4}CV^2$  (b)  $\frac{3}{4}CV^2$
- (c)  $\frac{1}{2}CV^2$  (d)  $2CV^2$ .
- A black body is at 727° C. It emits energy at a rate which is proportional to
  - (a)  $(1000)^4$
- (b)  $(1000)^2$
- (c)  $727)^4$
- (d)  $(727)^2$
- The resistance of an ammeter is 13  $\Omega$  and its scale is graduated for a current upto 100 amps. After an additional shunt has been connected to this ammeter it becomes possible to measure currents upto 750 amperes by this meter. The value of shunt-resistance is
  - (a)  $2\Omega$
- (b)  $0.2\Omega$  (c)  $2k\Omega$  (d)  $20\Omega$


- An engine has an efficiency of 1/6. When the temperature os sink is reduced by 62°C, its efficiency is doubled. Temperature of the source
  - (a) 37°C (b) 62°C (c) 99°C (d) 124°C
- The electric and magnetic field of an electromagnetic wave are
  - (a) in opposite phase and perpendicular to each other
  - (b) in opposite phase and parallel to each other
  - (c) in phase and perpendicular to each other
  - (d) in phase and parallel to each other.
- What is the value of inductance L for which the current is maximum in a series LCR circuit with  $C=10\,\mu F$  and  $\omega=1000s^{-1}$ 
  - (a) 1 mH
  - (b) cannot be calculated unless R is known
  - (c) 10mH
  - (d) 100mH
- 45. Nickel shows ferromagnetic property at room temperature. If the temperature is increased beyond Curie temperature, then it will show
  - (a) anti ferromagnetism
  - (b) no magnetic property
  - (c) diamagnetism
  - (d) paramagnetism
- 46. A vertical spring with force constant k is fixed on a table. A ball of mass m at a height h above the free upper end of the spring falls vertically on the spring so that the spring is compressed by a distance d. The net work done in the process
  - (a)  $mg(h+d) \frac{1}{2}kd^2$
  - (b)  $mg(h-d) \frac{1}{2}kd^2$
  - (c)  $mg(h-d) + \frac{1}{2}kd^2$
  - (d)  $mg(h+d) + \frac{1}{2}kd^2$
- 47. Under the influence of a uniform magnetic field a charged particle is moving in a circle of radius R with constant speed v. The time period of the motion
  - (a) depends on both R and v
  - (b) is independent of both R and v
  - (c) depends on R and not on v
  - (d) depends on v and not on R

- If the cold junction of a thermo-couple is kept at 0°C and the hot junction is kept at T°C then the relation between neutral temperature (T<sub>n</sub>) and temperature of inversion (T<sub>i</sub>) is

- (a)  $T_n = 2T_i$  (b)  $T_n = T_i T$ (c)  $T_n = T_i + T$  (d)  $T_n = T_i/2$ Assuming the sun to have a spherical outer 49. surface of radius r, radiating like a black body at temperature t°C, the power received by a unit surface, (normal to the incident rays) at a distance R from the centre of the sun is
  - (a)  $\frac{r^2\sigma(t+273)^4}{4\pi R^2}$  (b)  $\frac{16\pi^2r^2\sigma t^4}{R^2}$
  - (c)  $\frac{r^2\sigma(t+273)^4}{R^2}$  (d)  $\frac{4\pi r^2\sigma t^4}{R^2}$

where  $\sigma$  is the Stefan's constant.

In the energy band diagram of a material shown below, the open circles and filled circles denote holes and electrons respectively. The material is



- (a) an insulator
- (b) a metal
- (c) an n-type semiconductor
- (d) a p-type semiconductor

## CHEMISTRY

- Which one of the following on reduction with lithium aluminium hydride yields a secondary amine?
  - (a) Methyl isocyanide
  - (b) Acetamide
  - (c) Methyl cyanide
  - (d) Nitroethane.

- RNA and DNA are chiral molecules, their chirality is due to
  - (a) chiral bases
  - (b) chiral phosphate ester units
  - (c) D-sugar component
  - (d) L-sugar component.
- The Langmuir adsorption isotherm is deduced using the assumption
  - (a) the adsorption sites are equivalent in their ability to adsorb the particles
  - (b) the heat of adsorption varies with coverage
  - (c) the adsorbed molecules interact with each
  - (d) the adsorption takes place in multilayers.
- A weak acid, HA, has a  $K_a$  of  $1.00 \times 10^{-5}$ . If 0.100 mole of this acid disolved in one litre of water, the percentage of acid dissociated at equilbrium is closest to
  - (a) 1.00%
- (b) 99.9%
- (c) 0.100%
- (d) 99.0%
- The reaction of hydrogen and iodine monochloride is given as:

$$\mathbf{H}_{2(\mathbf{g})} + 2\mathbf{ICl}_{(\mathbf{g})} {\longrightarrow\!\!\!\!-\!\!\!\!-\!\!\!\!-\!\!\!\!-\!\!\!\!\!-} 2\mathbf{HCl}_{(\mathbf{g})} + \mathbf{I}_{2(\mathbf{g})}$$

The reaction is of first order with respect to  $H_{2(g)}$ and ICI<sub>(g)</sub>, following mechanisms were proposed. Mechanism A:

$$H_{2(g)} + 2ICl_{(g)} \longrightarrow 2HCl_{(g)} + I_{2(g)}$$

Mechanism B:

$$H_{2(g)} + ICl_{(g)} \longrightarrow HI_{(g)}$$
; slow

$$HI_{(g)} + ICl_{(g)} \longrightarrow HCl_{(g)} + I_{2(g)}$$
; fast

Which of the above mechanism(s) can be consistent with the given information about the reaction?

- (a) A and B both
- (b) neither A nor B
- (c) A only
- (d) B only
- The efficiency of a fuel cell is given by
  - (a)  $\frac{\Delta G}{\Delta S}$  (b)  $\frac{\Delta G}{\Delta H}$  (c)  $\frac{\Delta S}{\Delta G}$  (d)  $\frac{\Delta H}{\Delta G}$

- Consider the following reactions:
  - $H^{+}_{(aq)} + OH^{-}_{(aq)} = H_{2}O_{(l)},$   $\Delta H = -X_{1} \text{ kJ mol}^{-1}$
  - (ii)  $H_{2(g)} + \frac{1}{2}O_{2(g)} = H_2O(l)$ ,

 $\Delta H = -X_2 \, kJ \, mol^{-1}$ 

(iii)  $CO_{2(g)} + \overset{\checkmark}{H}_{2(g)} = CO_{(g)} + H_2O,$   $\Delta H = -X_3 \text{ kJ mol}^{-1}$ 

(iv)  $C_2H_{2(g)} + \frac{5}{2}O_{2(g)} = 2CO_2(g) + H_2O(l)$ 

 $\Delta H = -X_3 \, kJ \, mol^{-1}$ 

- Enthalpy of formation of  $H_2O_{(l)}$  is (a)  $+X_3$  kJ mol<sup>-1</sup> (b)  $-X_4$  kJ mol<sup>-1</sup> (c)  $+X_1$  kJ mol<sup>-1</sup> (d)  $-X_2$  kJ mol<sup>-1</sup>
- The correct order of increasing thermal stability of K<sub>2</sub>CO<sub>3</sub>, MgCO<sub>3</sub>, CaCO<sub>3</sub> and BeCO<sub>3</sub> is
  - (a)  $BeCO_3 < MgCO_3 < CaCO_3 < K_2CO_3$
  - (b)  $MgCO_3 < BeCO_3 < CaCO_3 < K_2CO_3$
  - (c)  $K_2CO_3 < MgCO_3 < CaCO_3 < BeCO_3$
  - (d)  $BeCO_3 < MgCO_3 < K_2CO_3 < CaCO_3$
- In which of the following pairs, the two species are iso-structure?
- Which of the following oxidation states are the most characteristic for lead and tin respectively?
  - (a) +2, +4
- (b) +4, +4
- (c) +2, +2
- (d) +4, +2
- Which of the following statements, about the advantage of roasting of sulphide ore before reduction is not true?
  - (a) The  $\Delta G_f^o$  of the sulphide is greater than those for CS<sub>2</sub> and H<sub>2</sub>S.
  - (b) The  $\Delta G_f^0$  is negative for roasting of sulphide ore to oxide.
  - Roasting of the sulphide to the oxide is thermodynamically feasible.
  - Carbon and hydrogen are suitable reducing agents for metal sulphides.
- 62. In a first-order reaction  $A \rightarrow B$ , if k is rate constant and inital concentration of the reactant A is 0.5 M, then the half-life is

- 63. Consider the following compounds.
  - (i) C<sub>6</sub>H<sub>5</sub>COCl

The correct decreasing order of their reactivity towards hydrolysis is

- (a) (i) > (ii) > (iii) > (iv)
- (b) (iv) > (ii) > (ii) > (iii)
- (c) (ii)>(iv)>(i)>(iii)
- (d) (ii) > (iv) > (iii) > (i)
- 64. Predict the product C obtained in the following reaction of butyne-1.

$$CH_3CH_2 - C \equiv CH + HCl \longrightarrow B \xrightarrow{HI} C$$

(a) 
$$CH_3 - CH_2 - CH_2 - C - H$$

$$\begin{array}{c} & I \\ | \\ \text{(b)} \quad \text{CH}_3 - \text{CH}_2 - \text{CH} - \text{CH}_2 \text{Cl} \end{array}$$

$$\begin{array}{ccc} \text{(d)} & \text{CH}_3 - \text{CH} - \text{CH}_2 \text{CH}_2 \text{I} \\ & & \mid & \text{Cl} \end{array}$$

- 65. Concentrated aqueous sulphuric acid is 98% H<sub>2</sub>SO<sub>4</sub> by mass and has a density of 1.80 g mL<sup>-1</sup>. Volume of acid required to make one litre of 0.1MH<sub>2</sub>SO<sub>4</sub> solution is
  - (a)  $16.65 \,\text{mL}$
- (b) 22.20 mL
- (c) 5.55 mL
- (d) 11.10mL
- 66. Reduction of aldehydes and ketones into hydrocarbons using zinc amalgam and conc. HCl is called
  - (a) Cope reduction
  - (b) Dow reduction
  - (c) Wolf-Kishner reduction
  - (d) Clemmensen redcution.

- 67. Given that bond energies of H H and Cl Cl are  $430 \text{ kJ mol}^{-1}$  and  $240 \text{ kJ mol}^{-1}$  respectively and  $\Delta H_f$  for HCl is  $90 \text{ kJ mol}^{-1}$ , bond enthalpy of HCl is
  - (a)  $380 \, kJ \, mol^{-1}$
- (b)  $425 \text{ kJ mol}^{-1}$
- (c)  $245 \text{ kJ mol}^{-1}$
- (d)  $290 \, \text{kJ} \, \text{mol}^{-1}$
- 68. Which of the following represents the correct order of the acidity in the given compounds?
  - (a) FCH<sub>2</sub>COOH > CH<sub>3</sub>COOH > BrCH<sub>2</sub>COOH > CICH<sub>2</sub>COOH
  - (b) BrCH<sub>2</sub>COOH>ClCH<sub>2</sub>COOH>FCH<sub>2</sub>COOH >CH<sub>3</sub>COOH
  - (c) FCH<sub>2</sub>COOH > CICH<sub>2</sub>COOH > BrCH<sub>2</sub>COOH > CH<sub>3</sub>COOH
  - (d) CH<sub>3</sub>COOH>BrCH<sub>2</sub>COOH>CICH<sub>2</sub>COOH> FCH<sub>2</sub>COOH
- 59. If threre is no rotation of plane polarised light by a compound in a specific solvent, though to be chiral, it may mean that
  - (a) the compound is certainly meso
  - (b) there is no compound in the solvent
  - (c) the compound may be a racemic mixture
  - (d) the compound is certainly a chiral.
- 70. CH<sub>3</sub> CHCl CH<sub>2</sub> CH<sub>3</sub> has a chiral centre. which one of the following represents its R-configuration?

- 71. Which of the following is water-soluble?
  - (a) Vitamin E
- (b) Vitamin K
- (c) Vitamin A
- (d) Vitamin B
- 72. Identify the correct order of the size of the following:
  - (a)  $Ca^{2+} < K^+ < Ar < Cl^- < S^{2-}$
  - (b)  $Ar < Ca^{2+} < K^+ < Cl^- < S^{2-}$
  - (c)  $Ca^{2+} < Ar < K^+ < Cl^- < S^{2-}$
  - (d)  $Ca^{2+} < K^+ < Ar < S^{2-} < Cl^-$
- 73. 0.5 molal aqueous solution of a weak acid (HX) is 20% ionised. If K<sub>f</sub> for water is 1.86 K kg mol–1,the lowering in freezing point of the solution is
  - (a) 0.56 K
- (b) 1.12 K
- (c)  $-0.56 \,\mathrm{K}$
- (d) -1.12 K

- Sulphide ores of metals are usually concentrated by froth flotation process. Which one of the following sulphide ores offer an exception and its concentrated by chemical leaching?
  - (a) Galena
- (b) Copper pyrite
- (c) Sphalerite
- (d) Argentite
- If 60% of a first order reaction was completed in 60 minutes, 50% of the same reaction would be completed in aproximately
  - (a) 45 minutes
- (b) 60 minutes
- (c) 40 minutes
- (d) 50 minutes

 $(\log 4 = 0.60, \log 5 = 0.69)$ 

- Which of the following will give a pair of enantiomorphs?
  - (a) [Cr(NH<sub>3</sub>)<sub>6</sub>][Co(CN)<sub>6</sub>]
  - (b)  $[Co(en)_2Cl_2]Cl$
  - (c)  $[Pt(NH_3)_4][PtCl_6]$
  - (d)  $[Co(NH_3)_4Cl_2]NO_2$ . (en=NH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)
- If NaCl is doped with  $10^{-4}$  mol \% of SrCl<sub>2</sub>, the concentration of cation vacancies will be  $(N_A = 6.02 \times 10^{23} \,\text{mol}^{-1})$ 
  - (a)  $6.02 \times 10^{16} \,\text{mol}^{-1}$  (b)  $6.02 \times 10^{17} \,\text{mol}^{-1}$
  - (c)  $6.02 \times 10^{14} \,\mathrm{mol^{-1}}$  (d)  $6.02 \times 10^{15} \,\mathrm{mol^{-1}}$
- The d electron configurations of Cr<sup>2+</sup>, Mn<sup>2+</sup>,  $Fe^{2+}$  and  $Ni^{2+}$  are  $3d^4$ ,  $3d^5$ ,  $3d^6$  and  $3d^8$ respectively. Which one of the following aqua complexes will exhibit the minimum paramagnetic behaviour?
  - (a)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$
- (b)  $[Ni(H_2O)_6]^{2+}$
- (c)  $[Cr(H_2^2O)_6]^{2+}$
- (d)  $[Mn(\tilde{H}_2O)_6]^{2+}$
- (At. No. Cr = 24, Mn = 25, Fe = 26, Ni = 28)
- In which of the following the hydration energy is higher than the lattice energy?
  - (a)  $MgSO_4$
- (b) RaSO<sub>4</sub>
- (c) SrSO<sub>4</sub>
- (d)  $BaSO_4$
- Which of the compounds with molecular formula  $C_5H_{10}$  yields acetone on ozonolysis?
  - (a) 3-methyl-1-butene
  - (b) cyclopentane
  - (c) 2-methyl-1-butene
  - (d) 2-methyl-2-butene.
- 81. With which of the following electronic configuration an atom has the lowest ionisation enthalpy?
  - (a)  $1s^2 2s^2 sp^3$
- (b)  $1s^2 2s^2 2p^5 3s^1$
- (c)  $1s^2 2s^2 2p^6$
- (d)  $1s^2 2s^2 2p^5$
- Which one of the following ionic species has the greatest proton affinity to form stable compound?
  - (a)  $NH_2^-$  (b)  $F^-$  (c)  $I^-$
- (d) HS-

The following equilibrium constants are given:

$$N_2 + 3H_2 \Longrightarrow 2NH_3; K_1$$

$$N_2 + O_2 \Longrightarrow 2NO; K_2$$

$$H_2 + \frac{1}{2}O_2 \longrightarrow H_2O; K_2$$

The equilibrium constant for the oxidation of NH<sub>3</sub> by oxygen to give NO is

- (a)  $\frac{K_2 K_3^2}{K_1}$  (b)  $\frac{K_2^2 K_3}{K_1}$
- (b)  $\frac{K_1K_2}{K_3}$
- (d)  $\frac{K_2K_3^3}{K_1}$
- Which one of the following ions is the most stable in aqueous solution?
  - (a) **V**<sup>3+</sup>
- (b) Ti<sup>3+</sup>
- (c)  $Mn^{3+}$
- (d) Cr<sup>3+</sup> (At.No. Ti = 22, V = 23, Cr = 24, Mn = 25)

The correct order of C -O bond length among

- $CO, CO_3^{2-}, CO_2$  is
- (a) CO < CO<sub>3</sub><sup>2-</sup> < CO<sub>2</sub> (b) CO<sub>3</sub><sup>2-</sup> < CO<sub>2</sub> < CO (c) CO < CO<sub>2</sub> < CO<sub>3</sub><sup>2-</sup>

- (d)  $CO_2 < C\tilde{O}_3^{2-} < \tilde{C}O$
- Which one of the following orders correctly represents the increasing acid strengths of the given acids?
  - (a) HOClO < HOCl < HOClO<sub>3</sub> < HOClO<sub>2</sub>
  - (b) HOClO<sub>2</sub> < HOClO<sub>3</sub> < HOClO < HOCl
  - (c) HOClO<sub>3</sub> < HOClO<sub>2</sub> < HOClO < HOCl
  - (d) HOCl<HOClO<HOClO<sub>2</sub><HOClO<sub>3</sub>
- Which one of the following on treatment with 50% aqueous sodium hydroxide yields the corresponding alcohol and acid?
  - (a) C<sub>6</sub>H<sub>5</sub>CHO
- (b) CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CHO

(c)  $CH_3 - C - CH_3$  (d)  $C_6H_5CH_2CHO$ 

- For (i) I-, (ii) Cl-, (iii) Br-, the increasing order of 88. nucleophilicity would be
  - (a)  $Cl^- < Br^- < I^-$
- (b)  $I^- < Cl^- < Br^-$
- (c)  $Br^- < Cl^- < I^-$
- (d)  $I^- < Br^- < Cl^-$
- Calculate the pOH of a solution at 25°C that contains  $1 \times 10^{-10}$  M of hydronium ions, i.e.  $H_2O^+$ .
  - (a) 4.000 (b) 9.0000(c) 1.000 (d) 7.000

- 90. Consider the following sets of quantum numbers:
  - n 1 m S (i) 3 0 0 +1/2
  - (ii) 2 2 1 +1/2
  - -23 (iii) 4 -1/2
  - (iv) 1 -1-1/2
  - +1/2(v) 3

Which of the following sets of quantum number is not possible?

- (a) (i), (ii), (iii) and (iv)(b) (ii), (iv) and (v)
- (c) (i) and (iii)
  - (d) (ii), (iii) and (iv)
- 91. The equilibrium constant of the reaction:

$$Cu_{(s)} + 2Ag^{+}_{(aq)} \longrightarrow Cu^{2+}_{(aq)} + 2Ag_{(s)};$$

 $E^{\circ} = 0.46 \text{ V}$  at 298 K is

- (a)  $2.0 \times 10^{10}$
- (b)  $4.0 \times 10^{10}$
- (c)  $4.0 \times 10^{15}$
- (d)  $2.4 \times 10^{10}$
- Which one of the following polymers is prepared by condensation polymerisation?
  - (a) Teflon
- (b) Natural rubber
- (c) Styrene
- (d) Nylon-66
- The order of decreasing reactivity towards an electrophilic reagent, for the following would be
  - (i) benzene
- (ii) toluene
- (iii) chlorobenzene
- (iv) phenol (a) (ii) > (iv) > (i) > (iii)
- (b) (iv) > (iii) > (ii) > (i)
- (c) (iv) > (ii) > (ii) > (iii)
- (d) (i) > (ii) > (iii) > (iv)
- Which of the following anions is present in the chain structure of silicates?
  - (a)  $(Si_2O_5^{2-})_n$
- (b)  $(SiO_3^{2-})_n$ (d)  $Si_2O_7^{6-}$

- An element, X has the following isotopic composition:

 $^{200}X:90\%$ 

199X: 8.0 %

The weighted average atomic mass of the naturally occuring element X is closest to

- (a) 201 amu
- (b) 202 amu
- (c) 199 amu
- (d) 200 amu
- The fraction of total volume occupied by the atoms present in a simple cube is

In the reaction:

$$\begin{array}{c} \text{CH}_3 \\ \mid \\ \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{O} - \text{CH}_2 - \text{CH}_3 + \text{HI} \xrightarrow{\quad \text{Heated} \quad} \end{array}$$

Which of the following compounds will be

(a) 
$$CH_3 - CH - CH_3 + CH_3CH_2OH$$
  
 $CH_3$ 

(b) 
$$CH_3 - CH - CH_2OH + CH_3CH_3$$
  
 $CH_3$ 

$$\begin{array}{c} CH_3\\ \mid\\ (c) \quad CH_3-CH-CH_2OH+CH_3-CH_2-I \end{array}$$

$$\begin{array}{c} \text{CH}_3\\ \mid\\ \text{(d)} \quad \text{CH}_3-\text{CH}-\text{CH}_2-\text{I}+\text{CH}_3\text{CH}_2\text{OH} \end{array}$$

- The product formed in Aldol condensation is
  - (a) a beta-hydroxy aldehyde or a beta-hydroxy
  - (b) an alpha-hydroxy aldehyde or ketone
  - (c) an alpha, beta unsaturated ester
  - (d) a beta-hydroxy acid
- The number of moles of KMnO<sub>4</sub> that will be needed to react with one mole of sulphite ion in acidic solution is
  - (a) 4/5
- (b) 2/5
- (c) 1
- 100. Identify the incorrect statement among the following:
  - (a) Lanthanoid contraction is the accumulation of successive shrinkages.
  - (b) As a result of lanthanoid contraction, the properties of 4d series of the transition elements have no similarities with the 5d series of elements.
  - (c) Shielding power of 4f electrons is quite weak.
  - (d) There is a decrease in the radii of the atoms or ions as one proceeds from La to Lu.

### **BIOLOGY**

- 101. ICBN stands for
  - (a) International Code of Botanical Nomenclature
  - (b) International congress of Biological Names
  - (c) Indian Code of Botanical Nomenclature
  - (d) Indian Congress of Biological Names.
- 102. Which pair of the following belongs to Basidiomycetes
  - (a) puffballs and Claviceps
  - (b) peziza and stink borns
  - (c) Morchella and mushrooms
  - (d) birds nest fungi and puffballs.
- 103. Biological organisation starts with
  - (a) cellular level (b) organismic level
  - (c) atomic level
  - (d) submicroscopic molecular level
- 104. Identify the odd combination of the habitat and the particular animal concerned.
  - (a) sunderbans
- Bengal Tiger
- (b) periyar
- Elephant
- (c) Rann of kutch
- wild Ass
- (d) Dachigam National park Snow Leopard.
- 105. If you suspect major deficiency of antibodies in a person to which of the following would you look for confirmatory evidence?
  - (a) serum globulins
  - (b) fibrinogen in the plasma
  - (c) haemocytes
- (d) serum albumins
- 106. A common test to find the genotype of a hybrid is by
  - (a) crossing of one  $F_2$  progeny with female parent
  - (b) studying the sexual behaviour of F<sub>1</sub>
  - (c) crossing of one F<sub>1</sub> progeny with male parent
  - (d) crossing of one  $F_2$  progeny with male parent.
- 107. The okazaki fragments in DNA chain growth
  - (a) polymerize in the 3' to 5' direction and forms replication fork
  - (b) prove semi-conservative natuire of DNA replication
  - (c) polymerize in teh 5' to 3' direction and explain 3' to 5' DNA replication
  - (d) result in transcription.

- 108. Which one of the following statement is correct?
  - (a) there is no evidence of the existence of gills during embryogensis of mammals
  - (b) all plant and animal cells are totipotent
  - (c) ontogeny repeats phylogeny
  - (d) stem cells are specialize cells.
- 109. In which one of the following the BOD (Biochemical Oxygen Demand) of sewage (s), distillery effluent (DE), paper mill effluent (PE) and sugar mill effluent (SE) have been arranged in ascending order?
  - (a) SE < PE < S < DE (b) PE < S < SE < DE
  - (c) S < DE < PE < SE (d) SE < S < PE < DE.
- 110. The concept of chemical evolution is based on
  - (a) interaction of water, air and clay under intense heat
  - (b) effect of solar radiation on chemicals
  - (c) possible orgin of life by combination of chemicals under suitable environmental conditions
  - (d) crystallization of chemicals.
- 111. Which one of the following is surrounded by a callose wall?
  - (a) male gamete (b) egg
  - (c) pollen grain
  - (d) microspore mother cell
- 112. Probiotics are
  - (a) cancer inducing microbes
  - (b) new kind of food allergens
  - (c) live microbial food supplement
  - (d) safe antibiotics
- 113. One of endangered species of Indian medicinal plants is that of
  - (a) ocimum
- (b) garlic
- (c) nepenthes
- (d) podophyllum
- 114. What is true about Nereis, scorpion, cockroach and silver fish?
  - (a) they all posses dorsal heart
  - (b) none of them is aquatic
  - (c) they all belong to the same phylum
  - (d) they all have jointed paired appendages
- 115. The living organisms can be unexceptionally distinguished from the non-living things on the basis of their ability for
  - (a) interaction with the environment and progressive evolution
  - (b) reproduction
  - (c) growth and movement
  - (d) responsiveness to touch.

- 116. A person who is one along hunger strike and is surviving only on water, will have.
  - (a) less amino acids in his urine
  - (b) more glucose in his blood
  - (c) less urea in his urine
  - (d) more sodium in his urine.
- 117. The length of DNA molecule greatly exceeds the dimensions of the nucleus in eukaryotic cells. How is this DNA accommodated?
  - (a) super-coiling in nucleosomes
  - (b) DNase digestion
  - (c) through elemination of repititive DNA
  - (d) deletion of non-essential genes.
- 118. A sequential expression of a set of human genes
  - (a) messenger RNA (b) DNA sequence
  - (c) ribosome
- (d) transfer RNA.
- 119. In a coal fired power plant electrostatic precipitators are installed to control emission of
  - (a)  $NO_{\times}$  (b) SPM (c) CO (d)  $SO_2$
- 120. Select the wrong statement fom the following.(a) Both chloroplasts and mitochondria have an internal compartment, the thylakoid
  - (b) Both chloroplasts and mitochondria contain DNA

space bounded by teh thylakoid membrane

- (c) The chloroplasts are generally much large than mitochondria
- (d) Both chloroplasts and mitochondria contain an inner and an outer membrane.
- 121. Geometric representation of age structure is a characteristic of
  - (a) population
- (b) landscape
- (c) ecosystem
- (d) biotic community.
- 122. During transcription, RNA polymerase holoenzyme binds to a gene promoter and assumes a saddle-like structure, what is it's DNA-binding sequence?
  - (a) AATT
- (b) CACC
- (c) TATA
- (d) TTAA
- 123. A plant requires magnesium for
  - (a) protein synthesis
  - (b) chlorophyll synthesis
  - (c) cell wall development
  - (d) holding cells together.
- 124. In the prothallus of a vascular cryptogam, the antherozoids and eggs mature a different times. As a result
  - (a) there is high degree of sterility
  - (b) one can conclude that the plant is apomictic

- (c) self-fertilization is prevented
- (d) there is no change in success rate of fertilization
- 125. Ergot of rye is caused by a species of
  - (a) uncimula
- (b) ustilago
- (c) claviceps
- (d) phytophthora.
- 126. The finches of Galapagos islands provide an evidence in favour of
  - (a) evolution due to mutation
  - (b) retrogressive evolution
  - (c) biogeographical evolution
  - (d) special creation.
- 127. Which one of the following is a slime mould?
  - (a) physarum
- (b) Thiobacillus
- (c) Anabaena
- (d) Rhizopus.
- 128. The population of an insect species shows an explosive increase in numbers during rainy season followed by its disappearance at the end of the season. What does this show?
  - (a) the food plants mature and die at the end of the rainy season
  - (b) its population growth curve is of J-type
  - (c) the population of its predators increases enormously
  - (d) S-shaped or sigmoid growth of this insect.
- 129. Which of the following is a flowering plant with nodules containing filamentous nitrogen-fixing micro-organism
  - (a) crotalaria juncea (b) cycas revoluta
  - (c) cicer arietinum
  - (d) casuarina equisetifolia
- 130. Which one of the following is being utilized as a source of biodiesel in the Indian countryside?
  - (a) betroot
- (b) sugarcane
- (c) pongamia
- (d) euphorbia
- 131. For a critical study of secondary growth in plants. which one of the following pairs is suitable?
  - (a) teak and pine
- (b) deodar and fern
- (c) wheat and maiden hair fern
- (d) sugarcane and sunflower.
- 132. Which one of the following is a fat-soluble vitamin and its related deficiency disease?
  - (a) Retinol
- Xerophtahalmia
- (b) Cobalamine
- Beri-beri
- (c) Calciferol
- Pellagra
  Scurvy.
- (d) A scorbic acid Scu

- 133. Two cells A and B are contiguous. Cell A has osmotic pressure 10 atm, turgor pressure 7 atm and diffusion pressure deficit 3 atm. Cell B has osmotic pressure 8 atm, turgor pressure 3 atm and diffusion pressure deficit 5 atm. The result will.
  - (a) no movement of water
  - (b) equilibrium between the two
  - (c) movement of water from cell A to B.
  - (d) movement of water from cell B to A.
- 134. In which one of the following preparations are your likely to come across cell junctions most frequently?
  - (a) thrombocytes (b) tendon
  - (c) hyaline cartilage (d) ciliated epithelium.
- 135. If you are asked to classify the various algae into district groups, which of the following characters you should choose?
  - (a) nature of stored food materials in the cell
  - (b) structural organization of thallus
  - (c) chemical composition of the cell wall
  - (d) types of pigments present in the cell.
- 136. About 98 percent of the mass of every living organism is composed of just six element including carbon, hydrogen, nitrogen, oxygen and
  - (a) sulphur and magnesium
  - (b) magnesium and sodium
  - (c) calcium and phosphorus
  - (d) phosphorus and sulphur.
- 137. Which one of the following statements is correct?
  - (a) Both Azotobacter and Rhizobium fix atmospheric nitrogen in root nodules of plants.
  - (b) Cyanobacteria such as Anabaena and Nostoc are important mobilizers of phosphates and for plant nutrition in soil
  - (c) At present it is not possible to grow maize without chemical fertilizers
  - (d) Extensive use of chemical fertilizers may lead to eutrophication of nearby water bodies.
- 138. In the hexaploid wheat, the haploid (n) and basic (x) numbers of chromosomes are
  - (a) n = 21 and x = 21 (b) n = 21 and x = 14
  - (c) n = 21 and x = 7 (d) n = 7 and x = 21.
- 39. Passage cells are thin walled cells found in
  - (a) phloem elements that serve as entry points for substance for transport ot other plant

- (b) testa of seeds to enable emergence of growing embryonic axis during seed germination
- (c) central region of style through which the pollen tube grows towards the ovary
- (d) endodermis of roots facilitating rapid transport of water from cortex to pericycle.
- 140. What is common between parrot, platypus and kangaroo?
  - (a) toothless jaws
  - (b) functional post-end tail
  - (c) ovoparity (d) homoiothermy
- 141. Bowman's glands are located in the
  - (a) anterior pituitary
  - (b) female reproductive system of cockroach
  - (c) olfactory epithelium of our nose
  - (d) proximal end of uriniferous tubles.
- 142. Which one of the following pairs is wrongly matched?
  - (a) yeast ethanol
  - (b) streptomycetes antibiotic
  - (c) coliforms vinegar
  - (d) methanogens gobar gas.
- 143. Which part of ovary in mammals acts as an endocrine gland after ovulation?
  - (a) stroma (b) germinal epithelium
  - (c) vetelline membrane
  - (d) Graafian follicle.
- 144. Opening of floral buds into flowers, is a type of
  - (a) autonomic movement of variation
  - (b) paratonic movement of growth
  - (c) autonomic movement of growth
  - (d) autonomic movement of locomotion.
- 145. Which one of the following is a matching pair of a body feature and the animal possessing it?
  - (a) ventral central Leech nervous system
  - (b) Pharyngeal gill slits Chamaeleon absent in embryo
  - (c) ventral heart Scorpion
  - (d) post-end tail Octopus.
- 146. All enzymes of TCA cycle are located in the mitochondrial matrix except one which is located in inner mitochondrial membranes in eukaryotes and in cytosol in prokaryotes. This enzyme is
  - (a) isocitrate dehydrogenase
  - (b) malate dehydrogenase
  - (c) succinate dehydrogenase
  - (d) lactate dehydrogenase.

- 147. The first acceptor of electrons from an excited chlorophyll molecule of photosystem II is
  - (a) iron-sulphur protein
  - (b) ferredoxin
  - (c) quinone
- (d) cytochrome
- 148. Differentiation of organs and tissues in a developing organism, is associated with
  - (a) differential expession of genes
  - (b) lethal mutations (c) deletion of genes
  - (d) developmental mutations.
- 149. Which one of the following statements about mycoplasma is wrong?
  - (a) They are pleomorphic
  - (b) They are sensitive to penicillin
  - (c) They cause diseases in plants
  - (d) They are also called PPLO.
- 150. Which one of the following pairs is mismatched?
  - (a) Apis indica honey
  - (b) kenia lacca lac
  - (c) Bombyx mori silk
  - (d) Pila globosa pearl
- 151. A person is having problems with calcium and phosphorus metabolism in his body. Which one of following glands may not be functioning properly?
  - (a) parotid
- (b) pancreas
- (c) thyroid
- (d) parathyroid
- 152. A genetically engineered micro-organism use successfully in bioremediation of oil spills is a species of
  - (a) Trichoderma
- (b) Xanthomonas
- (c) Bacillus
- (d) Pseudomonas.
- 153. In the leaves of C<sub>4</sub> plants, malic acid formation during CO<sub>2</sub> fixation occurs in the cells of
  - (a) bundle sheath
- (b) phloem
- (c) epidermis
- (d) mesophyll
- 154. Flagellated male gametes are present in all the three of which one of the following sets?
  - (a) Zygnema, Saprolegnia and Hydrilla
  - (b) fucus, Marsilea and Calotropis
  - (c) Riccia, Dryopteris and Cycas
  - (d) Anthoceros, Funaria and Spirogyra
- 155. In the mean and the median pertaining to a certain character of a population are of the same value, which of the follwing is most likely to occur
  - (a) a bi-modal distribution
  - (b) a T-shaped curve
  - (c) a skewed curve
  - (d) a normal distribution.

- 156. One gene-one enzyme relationship was established for the first time in
  - (a) Salmonella typhimurium
  - (b) Escherichia coli
  - (c) Diplocococcus pneumoniae
  - (d) Neurospora crassa.
- 157. Two genes R and Y are located very close on the chromosomal linkage map of maize plant. When RRYY and rryy genotypes are hybridized, the F<sub>2</sub> segrengation will show
  - (a) segregation in the expected 9:3:3:1 ratio
  - (b) segregation in 3:1 ratio
  - (c) higher number of the parental types
  - (d) higher number of the recombinant types.
- 158. In cloning of cattle, a fertillized egg is taken out of the mother's womb and
  - (a) in the eight cell stage, cells are separated and cultured untill small embryos are formed which are implanted into the womb of other cow
  - (b) in the eight cell stage the individual cells are separated under electrical field for further development in culture media.
  - (c) from this upto eight identical twins can be produced
  - (d) the egg is divided into 4 pairs of cells which are implanted into the womb of other cows
- 159. Ultrasound of how much frequency is beamed into human body for sonography?
  - (a) 15 30 MHZ
    - (b) 1 15 MHZ
  - (b) 45 70 MHZ
- (d) 30-45 MHZ.
- 160. Two plants can be conclusively said to belong to the same species if they.
  - (a) have more than 90 per cent similar genes
  - (b) look similar and possess identical secondary metabolites
  - (c) have same number of chromosomes
  - (d) can reproduce freely with each other and form seeds.
- 161. Inheritances of skin colour in humans is an example of
  - (a) point mutation
  - (b) polygenic inheritance
  - (c) codominance
  - (d) chromosomal aberration.
- 162. When two species of different genealogy come to resemble each other as a result of adaptation, the phenomenon is termed
  - (a) microevolution (b) co-evolution
  - (c) convergent evolution
  - (d) divergent evolution

- Increased asthmatics attacks in certain seasons are related to
  - (a) eating fruits preserved in tin containers
  - (b) inhalation of seasonal pollen
  - (c) low temperature
  - (d) hot and humid environment.
- Among the human ancestors the brain size was more that 1000 CC in
  - (a) Homo erectus (b) Ramapithecus
  - (c) Homo habilis
  - (d) Homo neanderthalensis
- Industrial melanism as observed in peppered moth proves that
  - (a) the melanic form of the moth has no selective advantage over lighter form in industrial area
  - (b) the lighter-form moth has no selective advantage either in polluted industrial area or non-polluted area
  - (c) melanism is a pollution-generated feature
  - (d) the true black melanic forms arise by a recurring random mutation.
- 166. A high density of elephant population in an area can result in
  - (a) intra specific competition
  - (b) inter specific competition
  - (c) predation on one another
  - (d) mutualism.
- 167. A drop of each of the following, is placed separately on four slides. Which of them will not coagulate?
  - (a) blood serum
  - (b) sample from the thoracic duct of lymphatic system
  - (c) Whole blood from pulmonary vein
  - (d) blood plasma.
- 168. Which one of the following ecosystem types has the highest annuial net primary productivity?
  - (a) tropical deciduous forest
  - (b) temperate evergreen forest
  - (c) temperate deciduous forest
  - (d) tropical rain forest.
- 169. In human body, which one of the following is anatomically correct?
  - (a) Collar bones 3 pairs
  - (b) Salivary glands 1 pairs
  - (c) Cranial nerves 10 pairs
  - (d) Floating ribs 2 pairs

- 170. Telomere repetitive DNA sequences control the function of eukaryotic chromosomes because they
  - (a) are RNA transcription initiator
  - (b) help chromosome pairing
  - (c) prevent chromosome loss
  - (d) act as replicons.
- 171. Spore diseminiation in some liverworts is aided by
  - (a) indusium
- (b) calyptra
- (c) peristome teeth (d) elaters.
- 172. Which one of the following elements is not an essential micronutrient for plant growth?
  - a) Zn (b) Cu (c) Ca (d) Mn
- 173. The overall goal of glycolysis, krebs cycle and the electron transport system is the formation of
  - (a) ATP in one large oxidation reaction
  - (b) sugars
- (c) nucleic acids
- (d) ATP in small stepwise units.
- 174. Adaptive radiation refers to
  - (a) evolution of different species from a common ancestor
  - (b) migration of members of a species to different geographical areas
  - (c) power of adaptation in an individual to a variety of environments
  - (d) adaptations due to geographical isolation.
- 175. Which one of the follwoing pairs, is not correctly matched?
  - (a) Gibberellic acid Leaf fall
  - (b) Cytokinin Cell wall elongation
  - (c) IAA Cell wall elongation
  - (d) Abscissic acid Stomatal closure.
- 176. Select the correct statement from the following?
  - (a) Fitness is the end result of the ability to adapt and gets selected by nature
  - (b) All mammals except whales and camels have seven cervical vertebrae
  - (c) Mutations are random and directional
  - (d) Darwinian variations are small and directionless.
- 177. Male gametes in angiosperms are formed by the division of
  - (a) generative cell (b) vegetative cell
  - (c) microspore mother cell
  - (d) microspore.
- 178. In pea plants, yellow seeds are dominant to green. If a heterozygous yellow seeded plant is crossed with a green seeded plant, what ratio of yellow and green seeded plants would you expect in F<sub>1</sub> generation?
  - (a) 9:1
- (b) 1:3
- (c) 3:1
- (d) 50:50

- 179. Which one of the follwing mammalian cells is not capable of metabolising glucose to carbondioxide aerobically?
  - (a) unstraited muscle cells
  - (b) liver cells
- (c) red blood cells
- (d) white blood
- 180. A human male produces sperms with the genotypes AB, Ab, aB, and ab pertaining to two diallelic characters in equal proportions. What is the corresponding genotype of this person?
  - (a) AaBB
- (b) AABb
- (c) AABB
- (d) AaBb.
- 181. What is common to whale, seal and shark?
  - (a) thick subcutaneous fat
  - (b) convergent evolution
  - (c) homoiothermy
  - (d) seasonal migration.
- 182. One of the important consequences of geographical isolation is
  - (a) preventing speciation
  - (b) speciation through reproductive isolation
  - (c) random creation of new species
  - (d) no change in the isolated fauna.
- 183. Lysozyme that is present in perspiration, saliva and tears, destroys
  - (a) certain types of bacteria
  - (b) all viruses
  - (c) most virus-infected cells
  - (d) certain fungi
- 184. In the human female, menstruation can be deferred by the administration of
  - (a) combination of FSH and LH
  - (b) combination of estrogen and progesterone
  - (c) FSH only
- (d) LH only
- 185. Which one of the following is not a constituent of cell membrane?
  - (a) glycolipids
- (b) proline
- (c) phospholipids (d) cholesterol.
- 186. Compared to a bull a bullock is docile because
  - higher levels of cortisone
  - lower levels of blood testosterone (b)
  - lower levels of adrenaline/ noradrenaline (c) in its blood
  - (d) higher levels of thyroxine.
- 187. During the transmission of nerve impulse through a nerve fibre, the potential on the inner side of the plasma membrane has which type of electric change?
  - (a) first positive, then negative and continue to be positive

- (b) first negative, then positive and continue to be positive.
- first positive, then negative and again back to positive
- first negative, then positive and again back to negative.
- 188. Which one of the following pairs of organisms are exotic species introduced in India?
  - lantana camara, water hyacinth
  - (b) water hyacinth, prosopis cinereria
  - nile perch, licus religiosa (c)
  - (d) ficus religiosa, lantana camara.
- 189. In gymnosperms, the pollen chamber represents
  - (a) a cavity in the ovule in which pollen grains are stored after pollination
  - an opening in the megagametophyte through which the pollen tube approaches the egg
  - (c) the microsporangium in which pollen grains develop
  - a cell in the pollen grain in which the sperms are formed.
- 190. Molecular basis of organ differentiation dpends on the modulation in transcription by
  - (a) ribosome
- (b) transcription factor
- (d) RNA polymerase. (c) anticodon
- 191. Which of the following is an example of negative feedback loop in humans?
  - secretion of tears after falling of sand particles into the eye.
  - salivation of mouth at the sight of delicious
  - secretion of sweat glands and constriction of skin blood vessels when it is too hot
  - constriction of skin blood vessels and contraction of skeletal mucles when it is too cold
- 192. The two polynucleotide chains in DNA are
  - (a) discontinuous (b) antiparallel
  - semiconservative(d) parallel.
- 193. Which of the following pairs are correctly matched?
  - Morphological features Animals
- Crocodile 4-chambered heart
- Sea urchin Parapodia (ii)
- Obelia Thecodont (iii)
  - Lemur Thecodont
    - (a) (ii), (iii) and (iv) (b) only (i) and (iv)
    - (c) only (i) and (ii) (d) (i), (iii) and (iv)

### 2007-16

- 194. Which one of the following pairs of structures distinguishes a nerve cell from other types of cells?
  - (a) vacuoles and fibres
  - (b) flagellum and modulloary sheath
  - (c) nucleus and mitochondria
  - (d) perikaryon and dendrites.
- 195. In maize, hybrid vigour is exploited by
  - (a) crossing of two inbred parental lines
  - (b) harvesting seeds from the most productive plants
  - inducing mutations
  - (d) bombarding the seeds with DNA.
- 196. Which one of the following is a viral disease of poultry?
  - (a) coryza
- (b) new castle disease
- (c) pasteurellosis
- (d) salmonellosis.

### CBSE-PMT - 2007 SCREENING SOLVED PAPER

- 197. The wavelength of light absorbed by Pr form of phytochrome is
  - (a) 680 nm
- (b) 720 nm
- (c) 620 nm
- (d) 640 nm
- 198. Which one of the following is not a bioindicator of water pollution?
  - (a) blood-worms
- (b) stone flies
- (c) sewage fungus (d) sludge-worms.
- 199. Feeling the tremors of an earthquake a scared resident of seventh floor of a multistoryed building starts climbing down the stairs rapidly. Which hormone initiated this action?
  - (a) adrenaline
- (b) glucagon
- (b) gastrin
- (d) thyroxine
- "Foolish seedling" disease of rice led to the 200. discovery of
  - (a) ABA
- (b) 2,4-D
- (c) IAA
- (d) GA